Stopain® Clinical Gel

Stopain® Clinical Gel is formulated with 10% menthol plus dimethylsulfone (MSM) and glucosamine. Stopain® Clinical gel works on contact to provide fast, long-lasting pain relief of sprains, strains, bruises, arthritis, muscle aches, joint and back pain.

Available Sizes:

- 12 x 3g Packet Shelf Box
- 4 fl. oz. Tube
- 16 fl. oz. Pump Bottle
- 1 gallon Pump Bottle

Product Benefits:

- Advanced formulas contain 10% active menthol plus MSM and glucosamine
- Penetration enhancers provide fast, long-lasting pain relief
- Sweat resistant formula
- Pharmaceutical grade skin conditioners
- Cutting edge polymer technology allows for a smoother, more consistent application
- No dyes or fragrances
- Can be used with kinesiology tape without interfering with adhesion







MEDICINAL INGREDIENTS

PURPOSE

Menthol (dl-Menthol): 4%, Menthol (l-Menthol): 6%

Topical Analgesic

NON-MEDICINAL INGREDIENTS acrylates/vinyl isodecanoate crosspolymer, dimethylsulfone (MSM), eucalyptus essential oil, glucosamine sulfate, hydroxypropylcellulose, PEG-8 dimethicone, pentylene glycol, peppermint oil, purified water, SD alcohol 39C, triethanolamine

3 YEAR EXPIRATION with stability testing completed on-site for all products

REPEATED INSULT PATCH TESTING performed by an independent lab to determine the dermal irritation and sensitization potential of the products showed no negative reactions.

CAUTIONS AND WARNINGS: Call a Poison Control Center immediately if overdose or accidental ingestion occurs. Consult a healthcare practitioner prior to use if you are pregnant or breastfeeding. Keep out of reach of children. Stop use and consult a health care practitioner if symptoms worsen, or re-occur within a few days.

KNOWN ADVERSE REACTIONS: Stop use and get medical help right away if you experience pain, swelling or blistering. Stop use if hypersensitivity/allergy, rashes and/or burning discomfort occur.

USES: Temporarily relieves aches and pains of muscles and joints associated with one or more of the following: simple backache, lumbago, strains and sprains (involving muscles, tendons and/or ligaments), and arthritis.

ADULTS AND CHILDREN 12 YEARS AND OLDER: For external use only. Application should be supervised by an adult. Apply thinly and evenly to affected area up to 3-4 times per day. Rub and/or massage into skin until solution vanishes. Avoid contact with the eyes and mucous membranes. Do not apply to wounds or damaged skin. Do not apply with external heat, such as electric heating pad, as this may result in excessive skin irritation or skin burn. Do not tightly bandage.





ADDITIONAL INFORMATION: Flammable. Do not use near a flame. Store at room temperature.